

# Automated Theorem Proving

Prof. Dr. Jasmin Blanchette, Lydia Kondylidou,  
Yiming Xu, PhD, and Tanguy Bozec  
based on exercises by Dr. Uwe Waldmann

Winter Term 2024/25

## Exercises 7: Resolution Continued

**Exercise 7.1:** Find a strict total ordering  $\succ$  on the ground atoms  $P(b), P(c), Q, R$  such that

$$P(b) \vee \neg P(c) \succ_C \neg P(b) \vee P(c) \quad (1)$$

$$P(b) \vee P(b) \vee P(b) \vee R \succ_C P(b) \vee R \vee R \quad (2)$$

$$\neg P(b) \vee Q \succ_C P(c) \vee R \quad (3)$$

**Exercise 7.2:** Consider the following formulas:

$$F_1 = \forall x (S(x) \rightarrow \exists y (R(x, y) \wedge P(y)))$$

$$F_2 = \forall x (P(x) \rightarrow Q(x))$$

$$F_3 = \exists x S(x)$$

$$G = \exists x \exists y (R(x, y) \wedge Q(y))$$

Use ordered resolution to prove that  $\{F_1, F_2, F_3\} \models G$ . You may choose the selection function and the ordering on atoms.

Hint: You will need some preprocessing.

**Exercise 7.3:** Let  $\Sigma = (\Omega, \Pi)$  be a signature with  $\Omega = \{b/0, f/1\}$  and  $\Pi = \{P/2, Q/1, R/2\}$ . Suppose that the atom ordering  $\succ$  compares ground atoms by comparing lexicographically first the predicate symbols ( $P \succ Q \succ R$ ), then the size of the first argument, then the size of the second argument (if present). If at least one of the two atoms to be compared is nonground,  $\succ$  compares only the predicate symbols.

Let  $N$  be the following set of clauses:

$$P(f(x), x) \vee R(b, b) \quad (1)$$

$$\neg P(b, x) \vee \neg P(x, b) \vee Q(x) \quad (2)$$

$$Q(f(b)) \vee \neg Q(b) \vee R(f(x), b) \quad (3)$$

$$Q(b) \vee \neg R(f(x), f(x)) \quad (4)$$

$$\neg Q(x) \vee R(x, x) \quad (5)$$

- (a) Which literals are strictly maximal in the clauses of  $N$ ?
- (b) Which literals are maximal in the clauses of  $N$ ?
- (c) Which  $Res_{sel}^>$ -inferences are possible if  $sel$  selects no literals? What are their conclusions?
- (d) Is there a  $Res_{sel}^>$ -inference between the clause

$$P(x, f(x)) \vee R(b, b) \quad (1')$$

and clause (2) if  $sel$  selects no literals? Justify your answer.

- (e) Define a selection function  $sel$  such that  $N$  is saturated under  $Res_{sel}^>$ .

**Exercise 7.4:** In Sect. 3.12 of the lecture notes, the inference rules for ground resolution with ordering restrictions (without selection functions) are given by

(Ground) Ordered Resolution:

$$\frac{D \vee A \quad C \vee \neg A}{D \vee C} \quad \text{if } A \succ L \text{ for all } L \text{ in } D \text{ and } \neg A \succeq L \text{ for all } L \text{ in } C.$$

(Ground) Ordered Factorization:

$$\frac{C \vee A \vee A}{C \vee A} \quad \text{if } A \succeq L \text{ for all } L \text{ in } C.$$

This calculus is sound and refutationally complete for sets of ground clauses.

Suppose that we replace the ordering restriction for the first inference rule by “if  $A \succ L$  for all  $L$  in  $D$  and  $A \succeq L$  for all  $L$  in  $C$ .”

- (a) Is the calculus with this modification still sound? If yes, give a short explanation; if no, give a counterexample.
- (b) Is the calculus with this modification still refutationally complete? If yes, give a short explanation; if no, give a counterexample.

**Exercise 7.5:** Determine all strict total orderings  $\succ$  on the atomic formulas  $P, Q, R, S$  such that the associated clause ordering  $\succ_c$  satisfies the properties (1)–(3) simultaneously:

$$P \vee Q \succ_c \neg Q \quad (1)$$

$$R \vee Q \succ_c \neg P \vee \neg P \quad (2)$$

$$\neg R \vee \neg R \succ_c S \quad (3)$$

**Exercise 7.6:** Let  $\Sigma = (\Omega, \Pi)$  be a signature with  $\Omega = \{b/0, f/1\}$  and  $\Pi = \{P/1, Q/1\}$ . Suppose that the atom ordering  $\succ$  compares ground atoms by comparing lexicographically first the size of the argument and then the predicate symbols ( $P \succ Q$ ). Let  $N$  be the following set of clauses:

$$\neg P(x) \vee P(f(x)) \quad (1)$$

$$\neg Q(f(b)) \vee P(f^3(b)) \quad (2)$$

$$Q(b) \vee Q(f(b)) \quad (3)$$

where  $f^0(b)$  stands for  $b$  and  $f^{i+1}(b)$  stands for  $f(f^i(b))$ .

(a) Sketch what the set  $G_\Sigma(N)$  of all ground instances of clauses in  $N$  looks like. How is it ordered w.r.t. the clause ordering  $\succ_c$ ?

(b) Construct the candidate interpretation  $I_{G_\Sigma(N)}^\succ$  of the set of all ground instances of clauses in  $N$ . Is it a model of  $G_\Sigma(N)$ ?

**Exercise 7.7:** Let  $\Sigma = (\Omega, \Pi)$  be a signature with  $\Omega = \{b/0, f/1\}$  and  $\Pi = \{P/1, Q/1\}$ . Suppose that the atom ordering  $\succ$  compares ground atoms by comparing lexicographically first the predicate symbols ( $P \succ Q$ ) and then the size of the argument. Let  $N$  be the following set of clauses:

$$\neg Q(y) \vee P(y) \quad (1)$$

$$Q(x) \vee Q(f(x)) \quad (2)$$

(a) Sketch what the set  $G_\Sigma(N)$  of all ground instances of clauses in  $N$  looks like. How is it ordered w.r.t. the clause ordering  $\succ_c$ ?

(b) Construct the candidate interpretation  $I_{G_\Sigma(N)}^\succ$  of the set of all ground instances of clauses in  $N$ .

**Exercise 7.8:** Let  $N$  be a set of ground clauses, and let  $\succ$  be a total and well-founded atom ordering. Prove or refute: If every clause in  $N$  is redundant w.r.t.  $N$ , then every clause in  $N$  is a tautology.

**Exercise 7.9 (\*)**: Give an example of two different first-order clauses  $F$  and  $G$  such that  $F$  entails  $G$  and  $G$  is not redundant w.r.t.  $\{F\}$ . If necessary, specify the atom ordering used by the redundancy criterion.

**Exercise 7.10 (\*)**: Give a clause  $C$  such that an “Ordered Resolution with Selection” inference is possible from  $C$  and  $C$  and the inference is not redundant w.r.t.  $\{C\}$ .