

# Automated Theorem Proving

Prof. Dr. Jasmin Blanchette, Yiming Xu, PhD,  
Lydia Kondylidou, and Tanguy Bozec  
based on exercises by Dr. Uwe Waldmann

Winter Term 2025/26

## Exercises 8: Semantic Tableaux

**Exercise 8.1:** Show unsatisfiability of the set of formulas

$$P \rightarrow (Q \rightarrow R) \quad (1)$$

$$P \rightarrow Q \quad (2)$$

$$P \wedge \neg R \quad (3)$$

by exhibiting a strict tableau.

**Exercise 8.2:** Check whether the following propositional formulas are valid or not using semantic tableaux. Give a brief explanation. Use exactly the expansion rules given in the lecture.

(a)  $(P \rightarrow Q) \rightarrow ((P \vee R) \rightarrow (Q \vee R))$

(b)  $(P \vee Q) \rightarrow (P \wedge Q)$

**Exercise 8.3:** Check whether the following propositional formulas are valid or not using semantic tableaux. Give a brief explanation. Use exactly the expansion rules given in the lecture.

(a)  $(P \rightarrow Q) \rightarrow ((Q \rightarrow R) \rightarrow (P \rightarrow R)).$

(b)  $(R \wedge (R \rightarrow P)) \rightarrow (P \wedge \neg Q).$

**Exercise 8.4:** Determine the satisfiability of the following set of ground formulas using the tableau calculus:

$$P(b) \wedge \neg P(d) \quad (1)$$

$$P(c) \vee (P(b) \wedge P(d)) \quad (2)$$

$$P(c) \rightarrow \neg(P(b) \vee P(d)) \quad (3)$$

Use exactly the expansion rules given in the lecture. State explicitly whether the set is satisfiable and give an explanation for that statement.

**Exercise 8.5:** Extend the tableau calculus to support the following connectives:

- The Sheffer stroke, denoted  $|$ , is a binary connective meaning “not both.” Thus,  $F | G$  is equivalent to  $\neg F \vee \neg G$ .
- The Peirce arrow, denoted  $\downarrow$ , is a binary connective meaning “neither nor.” Thus,  $F \downarrow G$  is equivalent to  $\neg F \wedge \neg G$ .

**Exercise 8.6:** Refute the following set of formulas using the tableau calculus with ground instantiation:

$$\forall x \exists y P(x, y) \quad (1)$$

$$\exists z \forall w \neg P(f(z), w) \quad (2)$$

**Exercise 8.7:** Refute the following set of formulas using the free-variable tableau calculus:

$$\forall x \exists y P(x, y) \quad (1)$$

$$\exists z \forall w \neg P(f(z), w) \quad (2)$$