

Automated Theorem Proving

Prof. Dr. Jasmin Blanchette, Yiming Xu, PhD,
Tanguy Bozec, and Lydia Kondylidou
based on exercises by Dr. Uwe Waldmann

Winter Term 2025/26

Exercises 7: Resolution Continued

Exercise 7.1: Find a strict total ordering $\succ$ on the ground atoms $P(b), P(c), Q, R$ such that

$$P(b) \vee \neg P(c) \succ_C \neg P(b) \vee P(c) \quad (1)$$

$$P(b) \vee P(b) \vee P(b) \vee R \succ_C P(b) \vee R \vee R \quad (2)$$

$$\neg P(b) \vee Q \succ_C P(c) \vee R \quad (3)$$

Exercise 7.2: Consider the following formulas:

$$F_1 = \forall x (S(x) \rightarrow \exists y (R(x, y) \wedge P(y)))$$

$$F_2 = \forall x (P(x) \rightarrow Q(x))$$

$$F_3 = \exists x S(x)$$

$$G = \exists x \exists y (R(x, y) \wedge Q(y))$$

Use ordered resolution to prove that $\{F_1, F_2, F_3\} \models G$. You may choose the selection function and the ordering on atoms.

Hint: You will need some preprocessing.

Exercise 7.3: Let $\Sigma = (\Omega, \Pi)$ be a signature with $\Omega = \{b/0, f/1\}$ and $\Pi = \{P/2, Q/1, R/2\}$. Suppose that the atom ordering $\succ$ compares ground atoms by comparing lexicographically first the predicate symbols ($P \succ Q \succ R$), then the size of the first argument, then the size of the second argument (if present). If at least one of the two atoms to be compared is nonground, $\succ$ compares only the predicate symbols.

Let N be the following set of clauses:

$$P(f(x), x) \vee R(b, b) \quad (1)$$

$$\neg P(b, x) \vee \neg P(x, b) \vee Q(x) \quad (2)$$

$$Q(f(b)) \vee \neg Q(b) \vee R(f(x), b) \quad (3)$$

$$Q(b) \vee \neg R(f(x), f(x)) \quad (4)$$

$$\neg Q(x) \vee R(x, x) \quad (5)$$

- (a) Which literals are strictly maximal in the clauses of N ?
- (b) Which literals are maximal in the clauses of N ?
- (c) Which $Res_{sel}^>$ -inferences are possible if sel selects no literals? What are their conclusions?
- (d) Is there a $Res_{sel}^>$ -inference between the clause

$$P(x, f(x)) \vee R(b, b) \quad (1')$$

and clause (2) if sel selects no literals? Justify your answer.

- (e) Define a selection function sel such that N is saturated under $Res_{sel}^>$.

Exercise 7.4: In Sect. 3.12 of the lecture notes, the inference rules for ground resolution with ordering restrictions (without selection functions) are given by

(Ground) Ordered Resolution:

$$\frac{D \vee A \quad C \vee \neg A}{D \vee C} \quad \text{if } A \succ L \text{ for all } L \text{ in } D \text{ and } \neg A \succeq L \text{ for all } L \text{ in } C.$$

(Ground) Ordered Factorization:

$$\frac{C \vee A \vee A}{C \vee A} \quad \text{if } A \succeq L \text{ for all } L \text{ in } C.$$

This calculus is sound and refutationally complete for sets of ground clauses.

Suppose that we replace the ordering restriction for the first inference rule by “if $A \succ L$ for all L in D and $A \succeq L$ for all L in C .”

- (a) Is the calculus with this modification still sound? If yes, give a short explanation; if no, give a counterexample.
- (b) Is the calculus with this modification still refutationally complete? If yes, give a short explanation; if no, give a counterexample.

Exercise 7.5: Determine all strict total orderings $\succ$ on the atomic formulas P, Q, R, S such that the associated clause ordering $\succ_c$ satisfies the properties (1)–(3) simultaneously:

$$P \vee Q \succ_c \neg Q \quad (1)$$

$$R \vee Q \succ_c \neg P \vee \neg P \quad (2)$$

$$\neg R \vee \neg R \succ_c S \quad (3)$$

Exercise 7.6: Let $\Sigma = (\Omega, \Pi)$ be a signature with $\Omega = \{b/0, f/1\}$ and $\Pi = \{P/1, Q/1\}$. Suppose that the atom ordering $\succ$ compares ground atoms by comparing lexicographically first the size of the argument and then the predicate symbols ($P \succ Q$). Let N be the following set of clauses:

$$\neg P(x) \vee P(f(x)) \quad (1)$$

$$\neg Q(f(b)) \vee P(f^3(b)) \quad (2)$$

$$Q(b) \vee Q(f(b)) \quad (3)$$

where $f^0(b)$ stands for b and $f^{i+1}(b)$ stands for $f(f^i(b))$.

(a) Sketch what the set $G_\Sigma(N)$ of all ground instances of clauses in N looks like. How is it ordered w.r.t. the clause ordering $\succ_c$?

(b) Construct the candidate interpretation $I_{G_\Sigma(N)}^\succ$ of the set of all ground instances of clauses in N . Is it a model of $G_\Sigma(N)$?

Exercise 7.7: Let $\Sigma = (\Omega, \Pi)$ be a signature with $\Omega = \{b/0, f/1\}$ and $\Pi = \{P/1, Q/1\}$. Suppose that the atom ordering $\succ$ compares ground atoms by comparing lexicographically first the predicate symbols ($P \succ Q$) and then the size of the argument. Let N be the following set of clauses:

$$\neg Q(y) \vee P(y) \quad (1)$$

$$Q(x) \vee Q(f(x)) \quad (2)$$

(a) Sketch what the set $G_\Sigma(N)$ of all ground instances of clauses in N looks like. How is it ordered w.r.t. the clause ordering $\succ_c$?

(b) Construct the candidate interpretation $I_{G_\Sigma(N)}^\succ$ of the set of all ground instances of clauses in N .

Exercise 7.8: Let N be a set of ground clauses, and let $\succ$ be a total and well-founded atom ordering. Prove or refute: If every clause in N is redundant w.r.t. N , then every clause in N is a tautology.

Exercise 7.9 (*): Give an example of two different first-order clauses F and G such that F entails G and G is not redundant w.r.t. $\{F\}$. If necessary, specify the atom ordering used by the redundancy criterion.

Exercise 7.10 (*): Give a clause C such that an “Ordered Resolution with Selection” inference is possible from C and C and the inference is not redundant w.r.t. $\{C\}$.